

# **Narrow-band, GeV gold ion beams from ultra-thin foils irradiated by intense sub-picosecond pulses**

P. Martin<sup>1</sup>, H. Ahmed<sup>2</sup>, A. Alejo<sup>3</sup>, M. Cercez<sup>4</sup>, D. Doria<sup>5</sup>, D. Gwynne<sup>1</sup>, F. Hanton<sup>1</sup>, J. Green<sup>2</sup>, A. Macchi<sup>7,8</sup>, D. Maclellan<sup>9</sup>, P. McKenna<sup>9</sup>, J. A. Ruiz<sup>6</sup>, M. Swantusch<sup>4</sup>, O. Willi<sup>4</sup>, S. Zhai<sup>1,10</sup>, M. Borghesi<sup>1</sup> and S. Kar<sup>1</sup>

<sup>1</sup> *Centre for Plasma Physics, School of Mathematics and Physics, Queen's University Belfast, BT7 1NN, UK*

<sup>2</sup> *Central Laser Facility, Rutherford Appleton Laboratory, Didcot, United Kingdom*

<sup>3</sup> *IGFAE, Universidade de Santiago de Compostela, Santiago de Compostela, Spain*

<sup>4</sup> *Heinrich-Heine-Universität, Düsseldorf, Germany*

<sup>5</sup> *ELI-NP, Magurele, Ilfov, Romania*

<sup>6</sup> *Universidad Politécnica de Madrid, Madrid, Spain*

<sup>7</sup> *Istituto Nazionale di Ottica, CNR, Pisa, Italy*

<sup>8</sup> *Department of Physics "E. Fermi", Pisa, Italy*

<sup>9</sup> *University of Strathclyde, Glasgow, United Kingdom*

<sup>10</sup> *Department of Mathematics and Physics, Shanghai Normal University, Shanghai, China*

Narrow energy band bunches of ions were produced from the interaction of intense ( $>10^{20}$  W/cm<sup>2</sup>), sub-picosecond-duration laser pulses with ultra-thin (15 nm) gold foils. These included the bulk target species, in particular the Au ions which are accelerated with spectral peaks centred at 1.5 GeV and with fluxes on the order of  $10^{12}$  particles per steradian, far surpassing Au ion fluxes reported by previous works by orders of magnitude [1,2]. 2D particle-in-cell simulations show a complex interplay between different acceleration mechanisms at different stages of interaction, suggesting the Au bunches stem from strong radiation pressure acceleration on a heavy-ion dominant plasma in the moments just before transparency, followed by an efficient acceleration due to transparency-enhanced mechanisms. We show that this effect is scalable to future multi-PW systems, where Au ion bunches at energies of several GeV are feasible.

[1] P. Wang, Z. Gong, S. G. Lee, Y. Shou, Y. Geng, C. Jeon, I. J. Kim, H.W. Lee, J. W. Yoon, J. H. Sung, S. K. Lee, D. Kong, J. Liu, Z. Mei, Z. Cao, Z. Pan, I. W. Choi, X. Yan, C. H. Nam, and W. Ma, *Physical Review X* **11**, 21049 (2021).

[2] F. H. Lindner, E. McCary, X. Jiao, T. M. Ostermayr, R. Roycroft, G. Tiwari, B. M. Hegelich, J. Schreiber, and P. G. Thirolf, *Plasma Physics and Controlled Fusion* **61**, 055002 (2019).